



Leukemia Inhibitory Factor Receptor (LIFR) Antibody


Catalogue No.: abx103730




Western blot analysis of recombinant Human LIFR.



Western blot analysis of Pig Skeletal muscle lysate, using Rabbit Anti-Human LIFR Antibody (1 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 0.2 µg/ml).



IHC-P analysis of Human Kidney Tissue, with DAB staining, using Rabbit Anti-Human LIFR Antibody (20 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 2 µg/ml).



IHC-P analysis of Human Placenta Tissue, with DAB staining, using Rabbit Anti-Human LIFR Antibody (20 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 2 µg/ml).

Polyclonal Antibody to Leukemia Inhibitory Factor Receptor (LIFR).

Target: Leukemia Inhibitory Factor Receptor (LIFR)

Clonality: Polyclonal

Reactivity: Human

Datasheet

Version: 2.0.0
Revision date: 13 Jul 2025



Tested Applications:	WB, IHC, IF/ICC
Host:	Rabbit
Recommended dilutions:	WB: 0.01-2 µg/ml, IHC: 5-20 µg/ml, IF/ICC: 5-20 µg/ml. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	abx067767 - Recombinant LIFR (Arg692-Ser833) expressed in E. coli
Form:	Liquid
Purification:	Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	P42702 (UniProt , ExPASy)
KEGG:	hsa:3977
String:	9606.ENSP00000263409
Buffer:	0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.
Specificity:	Cross-reacts with Pig LIFR.
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.