



X-Ray Repair Cross Complementing 6 (XRCC6) Antibody


Catalogue No.: abx103789




Western blot analysis of Human A549 cell lysate, using Mouse XRCC6 Antibody (3 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 0.2 µg/ml).




Western blot analysis of Human Hela cell lysate, using Rabbit Anti-Mouse XRCC6 Antibody (3 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 0.2 µg/ml).



Western blot analysis of the recombinant protein.




Western blot analysis of Human MCF7 cell lysate, using Rabbit Anti-Mouse XRCC6 Antibody (3 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 0.2 µg/ml).




Western blot analysis of the recombinant protein.

Datasheet

Version: 4.0.0
Revision date: 28 Aug 2025



IHC-P analysis of Mouse Heart Tissue, with DAB staining.



IHC-P analysis of Heart Tissue, with DAB staining.

Polyclonal Antibody to X-Ray Repair Cross Complementing 6 (XRCC6).

Target: X-Ray Repair Cross Complementing 6 (XRCC6)

Clonality: Polyclonal

Reactivity: Mouse

Tested Applications: WB, IHC, IF/ICC

Host: Rabbit

Recommended dilutions: WB: 0.01-2 µg/ml, IHC: 5-20 µg/ml, IF/ICC: 5-20 µg/ml. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: XRCC6 (Thr319-Val538)

Form: Liquid

Purification: Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.

Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

UniProt Primary AC: P23475 ([UniProt](#), [ExPASy](#))

Datasheet

Version: 4.0.0

Revision date: 28 Aug 2025



KEGG: mmu:14375

String: [10090.ENSMUSP00000068559](#)

Buffer: 0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only