



T-Lymphocyte Activation Antigen CD86 (CD86) Antibody


Catalogue No.: abx103992



Western blot analysis of recombinant Human CD86.



IHC-P analysis of Human Tonsil Tissue, with HE staining.



IHC-P analysis of Human Spleen Tissue, with HE staining.

Polyclonal Antibody to T-Lymphocyte Activation Antigen CD86 (CD86).

Target: T-Lymphocyte Activation Antigen CD86 (CD86)**Clonality:** Polyclonal**Reactivity:** Human**Tested Applications:** WB, IHC, IF/ICC**Host:** Rabbit**Recommended dilutions:** WB: 0.01-2 µg/ml, IHC: 5-20 µg/ml, IF/ICC: 5-20 µg/ml. Optimal dilutions/concentrations should be determined by the end user.**Conjugation:** Unconjugated

Datasheet

Version: 5.0.0
Revision date: 01 May 2025



Immunogen:	LAB7-2 (Gln46-Cys282)
Form:	Liquid
Purification:	Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	P42081 (UniProt , ExPASy)
KEGG:	hsa:942
String:	9606.ENSP00000332049
Buffer:	0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only