



Pulmonary Activation Regulated Chemokine (PARC) Antibody


Catalogue No.: abx104139




Western blot analysis of recombinant Human PARC.




Western blot analysis of Human Leukocyte lysate, using Rabbit Anti-Human PARC Antibody (5 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 0.2 µg/ml).



Western blot analysis of Human Lung lysate, using Rabbit Anti-Human PARC Antibody (5 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 0.2 µg/ml).



Western blot analysis of Mouse Lymphocyte lysate, using Rabbit Anti-Human PARC Antibody (5 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 0.2 µg/ml).



IHC-P analysis of Lung tissue, with DAB staining.

Datasheet

Version: 4.0.0
Revision date: 28 Feb 2025



Polyclonal Antibody to CCL18.

Target:	Pulmonary Activation Regulated Chemokine (PARC)
Clonality:	Polyclonal
Reactivity:	Human
Tested Applications:	WB, IHC, IF/ICC
Host:	Rabbit
Recommended dilutions:	WB: 0.01-2 µg/ml, IHC: 5-20 µg/ml, IF/ICC: 5-20 µg/ml. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	abx068791 - Recombinant PARC (Ala21-Ala89) expressed in E. coli
Form:	Liquid
Purification:	Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	P55774 (UniProt , ExPASy)
KEGG:	hsa:6362
String:	9606.ENSP00000479955
Buffer:	0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.