



Transaldolase (TALDO1) Antibody

Catalogue No.:abx005186



Western blot analysis of various lysates using TALDO1 Antibody at 1/1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1/10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Exposure time: 30s.



Immunofluorescence analysis of HeLa cells using TALDO1 Antibody. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) at 1/500 dilution. Blue: DAPI for nuclear staining.

TALDO1 Antibody is a Rabbit Polyclonal antibody against TALDO1. Transaldolase 1 is a key enzyme of the nonoxidative pentose phosphate pathway providing ribose-5-phosphate for nucleic acid synthesis and NADPH for lipid biosynthesis. This pathway can also maintain glutathione at a reduced state and thus protect sulfhydryl groups and cellular integrity from oxygen radicals. The functional gene of transaldolase 1 is located on chromosome 11 and a pseudogene is identified on chromosome 1 but there are conflicting map locations. The second and third exon of this gene were developed by insertion of a retrotransposable element. This gene is thought to be involved in multiple sclerosis.

Target: Transaldolase (TALDO1)

Clonality: Polyclonal

Reactivity: Human, Mouse

Tested Applications: ELISA, WB, IF/ICC

Host: Rabbit

 $\textbf{Recommended dilutions:} \ \ \text{ELISA:} \ \ 1 \ \mu g/ml, \ WB: \ 1/500 - 1/2000, \ IF/ICC: \ 1/10 - 1/100. \ \ Optimal \ dilutions/concentrations \ should$

be determined by the end user.

Conjugation: Unconjugated

Immunogen: Recombinant protein corresponding to TALDO1. The exact sequence is proprietary.

Isotype: IgG

Datasheet

Version: 5.0.0 Revision date: 14 Sep 2025



Form: Liquid

Purification: Purified by affinity chromatography.

Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

UniProt Primary AC: P37837 (<u>UniProt</u>, <u>ExPASy</u>)

Gene Symbol: TALDO1

GeneID: <u>6888</u>

NCBI Accession: NP_006746.1

KEGG: hsa:6888

String: <u>9606.ENSP00000321259</u>

Molecular Weight: Calculated MW: 38 kDa

Observed MW: 38 kDa

Buffer: PBS, pH 7.3, containing 0.02% sodium azide, 50% glycerol.

Concentration: > 0.2 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC,

THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL

CONSUMPTION.