



## Serine/threonine-Protein Kinase Tousled-Like 2 (TLK2) Antibody

Catalogue No.:abx005192



Western blot analysis of various lysates using [KO Validated] TLK2 Antibody at 1/1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1/10000 dilution. Lysates/proteins: 25  $\mu$ g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Exposure time: 90s.



Western blot analysis of lysates from wild type (WT) and TLK2 knockout (KO) NIH/3T3 cells, using [KO Validated] TLK2 Antibody at 1/500 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1/10000 dilution. Lysates/proteins: 25 μg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Exposure time: 180s.

TLK2 Antibody is a Rabbit Polyclonal antibody against TLK2. This gene encodes a nuclear serine/threonine kinase that was first identified in Arabidopsis. The encoded protein is thought to function in the regulation of chromatin assembly in the S phase of the cell cycle by regulating the levels of a histone H3/H4 chaperone. This protein is associated with double-strand break repair of DNA damage caused by radiation. Pseudogenes of this gene are present on chromosomes 10 and 17. Alternate splicing results in multiple transcript variants.

Target: Serine/threonine-Protein Kinase Tousled-Like 2 (TLK2)

Clonality: Polyclonal

Reactivity: Human, Mouse

Tested Applications: ELISA, WB

Host: Rabbit

Recommended dilutions: ELISA: 1 µg/ml, WB: 1/500 - 1/2000. Optimal dilutions/concentrations should be determined by the

end user.

Conjugation: Unconjugated

**Immunogen:** Recombinant protein corresponding to TLK2. The exact sequence is proprietary.

**Isotype:** IgG

1 of 2

## **Datasheet**

Version: 5.0.0 Revision date: 02 Aug 2025



Form: Liquid

**Purification:** Purified by affinity chromatography.

**Storage:** Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

UniProt Primary AC: Q86UE8 (<u>UniProt</u>, <u>ExPASy</u>)

Gene Symbol: TLK2

GeneID: <u>11011</u>

NCBI Accession: NP\_006843.2

Molecular Weight: Calculated MW: 88 kDa

Observed MW: 87 kDa

**Buffer:** PBS, pH 7.3, containing 0.02% sodium azide, 50% glycerol.

**Concentration:** > 0.2 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC,

THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL

CONSUMPTION.