



Translocase of Outer Mitochondrial Membrane 20 (TOMM20) Antibody


Catalogue No.: abx005196




Immunohistochemistry of paraffin-embedded Human colon carcinoma using TOM20 Antibody (1/100 dilution, 40x lens).




Immunohistochemistry of paraffin-embedded Human colon using TOM20 Antibody (1/100 dilution, 40x lens).



Immunohistochemistry of paraffin-embedded Human lung cancer using TOM20 Antibody (1/100 dilution, 40x lens).



Immunohistochemistry of paraffin-embedded Mouse kidney using TOM20 Antibody (1/100 dilution, 40x lens).




Immunohistochemistry of paraffin-embedded Rat heart using TOM20 Antibody (1/100 dilution, 40x lens).


Datasheet

Version: 4.0.0


Revision date: 19 Sep 2025



Confocal immunofluorescence analysis of U2OS cells using TOM20 Antibody (1/100 dilution). Blue: DAPI for nuclear staining.



Immunofluorescence analysis of C6 cells using TOM20 Antibody (1/100 dilution). Blue: DAPI for nuclear staining.



Immunofluorescence analysis of NIH/3T3 using TOM20 Antibody (1/100 dilution). Blue: DAPI for nuclear staining.

TOMM20 Antibody is a Rabbit Polyclonal antibody against TOMM20.

Target: Translocase of Outer Mitochondrial Membrane 20 (TOMM20)

Clonality: Polyclonal

Reactivity: Human, Mouse, Rat

Tested Applications: IHC, IF/ICC

Host: Rabbit

Recommended dilutions: IHC-P: 1/50 - 1/200, IF/ICC: 1/50 - 1/200. Not tested in IHC-F. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: Recombinant protein corresponding to human TOM20

Datasheet

Version: 4.0.0

Revision date: 19 Sep 2025



Isotype:	IgG
Form:	Liquid
Purification:	Purified by affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	Q15388 (UniProt , ExPASy)
Gene Symbol:	TOMM20
GeneID:	9804
NCBI Accession:	NP_055580.1
KEGG:	hsa:9804
String:	9606.ENSP00000355566
Buffer:	PBS, pH 7.3, containing 0.02% sodium azide, 50% glycerol.
Concentration:	1 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.