



Cytotoxic T-Lymphocyte Associated Antigen 4 (CTLA4) Antibody

Catalogue No.: abx104332



Western blot analysis of recombinant Rat CTLA4 Protein.



IHC-P analysis of Rat Kidney Tissue, with DAB staining, using Rabbit Anti-Rat CTLA4 Antibody (10 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 2 µg/ml).

Polyclonal Antibody to Cytotoxic T-Lymphocyte Associated Antigen 4 (CTLA4).

Target: Cytotoxic T-Lymphocyte Associated Antigen 4 (CTLA4)

Clonality: Polyclonal

Reactivity: Rat

Tested Applications: WB, IHC, IF/ICC

Host: Rabbit

Recommended dilutions: WB: 0.01-2 µg/ml, IHC: 5-20 µg/ml, IF/ICC: 5-20 µg/ml. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: [abx066269](#) - Recombinant CTLA4 (Thr41-Lys192) expressed in E. coli

Form: Liquid

Purification: Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.

Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

Datasheet

Version: 2.0.0
Revision date: 13 Jun 2025



UniProt Primary AC: Q62859 ([UniProt](#), [ExPASy](#))

Gene Symbol: CTLA4

GeneID: [63835](#)

KEGG: rno:63835

Buffer: 0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.

Specificity: Cross-reacts with Mouse CTLA4.

Concentration: 0.85 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only