



Transforming Growth Factor Beta Induced Protein (TGF β I) Antibody


Catalogue No.: abx104492



Western blot analysis of the recombinant protein.



IHC-P analysis of Human Stomach Tissue, with DAB staining.



IHC-P analysis of Liver tissue, with DAB staining.

Polyclonal Antibody to Transforming Growth Factor Beta Induced Protein (TGF β I).

Target: Transforming Growth Factor Beta Induced Protein (TGF β I)

Clonality: Polyclonal

Reactivity: Human

Tested Applications: WB, IHC, IF/ICC

Host: Rabbit

Recommended dilutions: WB: 0.01-2 μ g/ml, IHC: 5-20 μ g/ml, IF/ICC: 5-20 μ g/ml. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Datasheet

Version: 2.0.0

Revision date: 27 Sep 2025



Immunogen:	TGFb1 (Gly423-Leu632)
Form:	Liquid
Purification:	Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	Q15582 (UniProt , ExPASy)
KEGG:	hsa:7045
String:	9606.ENSP00000416330
Buffer:	0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only