



Peroxisome Proliferator Activated Receptor Alpha (PPARα) Antibody


Catalogue No.: abx104506




Western blot analysis of Pig Skeletal Muscle Tissue.




Western blot analysis of recombinant Human PPARα.



IHC-P analysis of Human Liver Tissue, with DAB staining.



IHC-P analysis of Human Liver Cancer Tissue, with DAB staining.



IHC-P analysis of Human Breast Cancer Tissue, with DAB staining.

Datasheet

Version: 1.0.0
Revision date: 11 Oct 2025



Polyclonal Antibody to Peroxisome Proliferator Activated Receptor Alpha (PPARA).

Target:	Peroxisome Proliferator Activated Receptor Alpha (PPARα)
Clonality:	Polyclonal
Reactivity:	Human
Tested Applications:	WB, IHC, IF/ICC
Host:	Rabbit
Recommended dilutions:	WB: 0.01-2 µg/ml, IHC: 5-20 µg/ml, IF/ICC: 5-20 µg/ml. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	PPARα (Pro61-Ala170)
Form:	Liquid
Purification:	Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	Q07869 (UniProt , ExPASy)
KEGG:	hsa:5465
String:	9606.ENSP00000385523
Buffer:	0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.
Concentration:	0.91 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.