



Peroxisome Proliferator Activated Receptor Gamma (PPARγ) Antibody


Catalogue No.: abx104512



Western blot analysis of recombinant Mouse PPARγ protein.



Western blot analysis of Rat heart lysate using PPARγ Antibody (1 μg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 0.2 μg/ml).



IHC-P analysis of paraffin embedded Mouse cardiac muscle tissue using PPARγ antibody (20 μg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 2 μg/ml).

Polyclonal Antibody to Peroxisome Proliferator Activated Receptor Gamma (PPARγ).

Target: Peroxisome Proliferator Activated Receptor Gamma (PPARγ)

Clonality: Polyclonal

Reactivity: Mouse

Tested Applications: WB, IHC, IF/ICC

Host: Rabbit

Recommended dilutions: WB: 0.01-2 μg/ml, IHC: 5-20 μg/ml, IF/ICC: 5-20 μg/ml. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Datasheet

Version: 5.0.0
Revision date: 24 May 2025



Immunogen:	abx068500 - Recombinant PPAR γ (Phe149-Ser273) expressed in E. coli.
Form:	Liquid
Purification:	Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	P37238 (UniProt , ExPASy)
KEGG:	mmu:19016
String:	10090.ENSMUSP00000000450
Buffer:	0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.
Specificity:	Cross-reacts with Human and Rat PPAR γ .
Concentration:	0.71 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.