



CD70 Antigen / Tumor Necrosis Factor Ligand Superfamily Member 7 (CD70) Antibody

Catalogue No.: abx104601



Western blot analysis of the recombinant protein.



IHC-P analysis of Ovary tissue, with DAB staining.

Polyclonal Antibody to CD70 (CD70).

Target: CD70 Antigen / Tumor Necrosis Factor Ligand Superfamily Member 7 (CD70)**Research Area:** Immunology**Clonality:** Polyclonal**Reactivity:** Human**Tested Applications:** WB, IHC, IF/ICC**Host:** Rabbit**Recommended dilutions:** WB: 0.01-2 µg/ml, IHC: 5-20 µg/ml, IF/ICC: 5-20 µg/ml. Optimal dilutions/concentrations should be determined by the end user.**Conjugation:** Unconjugated**Immunogen:** CD70 (Gln47-Val191)**Form:** Liquid

Datasheet

Version: 5.0.0
Revision date: 29 Aug 2025



Purification:	Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	P32970 (UniProt , ExPASy)
Gene Symbol:	CD70
GeneID:	970
KEGG:	hsa:970
String:	9606.ENSP00000245903
Buffer:	0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only