



Tumor Necrosis Factor Ligand Superfamily Member 14 / LIGHT (TNFSF14) Antibody


Catalogue No.:abx104634



Western blot analysis of the recombinant protein.



IHC-P analysis of Rat Intestine Tissue, with DAB staining.



IHC-P analysis of kidney tissue, with DAB staining.

Polyclonal Antibody to Tumor Necrosis Factor Ligand Superfamily, Member 14 (TNFSF14).

Tumor Necrosis Factor Ligand Superfamily Member 14 / LIGHT (TNFSF14)

Clonality: Polyclonal

Reactivity: Rat

Tested Applications: WB, IHC, IF/ICC

Host: Rabbit

Recommended dilutions: WB: 0.01-2 µg/ml, IHC: 5-20 µg/ml, IF/ICC: 5-20 µg/ml. Optimal dilutions/concentrations should be

determined by the end user.

Datasheet

Version: 3.0.0 Revision date: 14 Jul 2025



Conjugation: Unconjugated

Immunogen: TNFSF14 (Arg58-Ile239)

Form: Liquid

Purification: Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.

Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

Buffer: 0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC,

THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL

CONSUMPTION.