



T-Cell Immunoreceptor with Ig and ITIM Domains (TIGIT) Antibody


Catalogue No.: abx104647




Western blot analysis of recombinant Human TIGIT.




IHC-P analysis of Human Thyroid Cancer Tissue, with DAB staining.



IHC-P analysis of Human Stomach Tissue, with DAB staining.



IHC-P analysis of Human Esophagus Cancer Tissue, with DAB staining.



IHC-P analysis of Human Kidney Cancer Tissue, with DAB staining.

Datasheet

Version: 6.0.0
Revision date: 18 Mar 2025



Polyclonal Antibody to T-Cell Immunoreceptor with Ig and ITIM Domains (TIGIT).

Target:	T-Cell Immunoreceptor with Ig and ITIM Domains (TIGIT)
Clonality:	Polyclonal
Reactivity:	Human
Tested Applications:	WB, IHC, IF/ICC
Host:	Rabbit
Recommended dilutions:	WB: 0.01-2 µg/ml, IHC: 5-20 µg/ml, IF/ICC: 5-20 µg/ml. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	abx069243 - Recombinant TIGIT (Gly21-Leu234) expressed in E. coli
Form:	Liquid
Purification:	Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	Q495A1 (UniProt , ExPASy)
Gene Symbol:	TIGIT
GeneID:	201633
OMIM:	612859
HGNC:	26838
KEGG:	hsa:201633
Ensembl:	ENSG00000181847
String:	9606.ENSP00000419085
Buffer:	0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.
Concentration:	0.7 mg/ml

Datasheet

Version: 6.0.0

Revision date: 18 Mar 2025



Note:

THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only