



Tumor Necrosis Factor Ligand Superfamily Member 10 (TNFSF10) Antibody

Catalogue No.:abx104661




Western blot analysis of Human MCF7 Cells.




Western blot analysis of recombinant Rat TNFSF10.



IHC-P analysis of Rat Testis Tissue, with DAB staining.



IHC-P analysis of Rat Ovary Tissue, with DAB staining.




IHC-P analysis of Rat Kidney Tissue, with DAB staining.

Datasheet

Version: 4.0.0 Revision date: 06 Sep 2025





IHC-P analysis of Rat Stomach Tissue, with DAB staining.

Polyclonal Antibody to Tumor Necrosis Factor Ligand Superfamily Member 10 (TNFSF10).

Target: Tumor Necrosis Factor Ligand Superfamily Member 10 (TNFSF10)

Research Area: Apoptosis, Tumor Immunity

Clonality: Polyclonal

Reactivity: Rat

Tested Applications: WB, IHC, IF/ICC

Host: Rabbit

Recommended dilutions: WB: 0.01-2 μg/ml, IHC: 5-20 μg/ml, IF/ICC: 5-20 μg/ml. Optimal dilutions/concentrations should be

determined by the end user.

Conjugation: Unconjugated

Immunogen: TNFSF10 (Pro115-Ala287)

Form: Liquid

Purification: Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.

Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

UniProt Primary AC: A0A8I5ZY27 (UniProt, ExPASy)

Gene Symbol: TNFSF10

Buffer: 0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC,

THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL

CONSUMPTION.