



Protein Disulfide Isomerase A6 (PDIA6) Antibody


Catalogue No.:abx005339



Western blot analysis of various lysates using PDIA6 Antibody at 1/1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1/10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Exposure time: 60s.



Immunohistochemistry analysis of paraffin-embedded Rat brain using PDIA6 Antibody at dilution of 1/100 (40x lens). Microwave antigen retrieval performed in 0.01 M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse brain using PDIA6 Antibody at dilution of 1/100 (40x lens). Microwave antigen retrieval performed in 0.01 M PBS Buffer (pH 7.2) prior to IHC staining.

PDIA6 Antibody is a Rabbit Polyclonal antibody against PDIA6. Protein disulfide isomerases (EC 5.3.4.1), such as PDIA6, are endoplasmic reticulum (ER) resident proteins that catalyze formation, reduction, and isomerization of disulfide bonds in proteins and are thought to play a role in folding of disulfide-bonded proteins (Hayano and Kikuchi, 1995.

Target: Protein Disulfide Isomerase A6 (PDIA6)

Clonality: Polyclonal

Reactivity: Human, Mouse, Rat

Tested Applications: ELISA, WB, IHC

Host: Rabbit

Recommended dilutions: ELISA: 1 µg/ml, WB: 1/500 - 1/2000, IHC-P: 1/50 - 1/200. Not tested in IHC-F. Optimal

dilutions/concentrations should be determined by the end user.

Datasheet

Version: 11.0.0 Revision date: 13 Aug 2025



Conjugation: Unconjugated

Immunogen: Recombinant protein corresponding to PDIA6. The exact sequence is proprietary.

Isotype: IgG

Form: Liquid

Purification: Purified by affinity chromatography.

Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

UniProt Primary AC: Q15084 (<u>UniProt</u>, <u>ExPASy</u>)

Gene Symbol: PDIA6

GenelD: <u>10130</u>

NCBI Accession: NP_005733.1

KEGG: hsa:10130

String: <u>9606.ENSP00000385385</u>

Molecular Weight: Calculated MW: 48 kDa

Observed MW: 48 kDa

Buffer: PBS, pH 7.3, containing 0.02% sodium azide, 50% glycerol.

Concentration: > 0.2 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC,

THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL

CONSUMPTION.