



Histone H3K9me1 Antibody


Catalogue No.: abx000005




Western blot analysis of various lysates using MonoMethyl-Histone H3-K9 Antibody at 1/1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1/10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Exposure time: 5s.




Dot-blot analysis of all sorts of methylation peptides using MonoMethyl-Histone H3-K9 antibody.




Immunohistochemistry analysis of paraffin-embedded Human colon using MonoMethyl-Histone H3-K9 Antibody at dilution of 1/50 (40x lens). High pressure antigen retrieval performed in 0.01 M Citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Mouse lung using MonoMethyl-Histone H3-K9 Antibody at dilution of 1/50 (40x lens). High pressure antigen retrieval performed in 0.01 M Citrate buffer (pH 6.0) prior to IHC staining.




Immunofluorescence analysis of 293T cells using MonoMethyl-Histone H3-K9 Antibody. Blue: DAPI for nuclear staining.




Immunofluorescence analysis of C6 cells using MonoMethyl-Histone H3-K9 Antibody at dilution of 1/100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) at 1/500 dilution. Blue: DAPI for nuclear staining.




Immunofluorescence analysis of HeLa cells using MonoMethyl-Histone H3-K9 Antibody at dilution of 1/100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) at 1/500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of NIH/3T3 cells using MonoMethyl-Histone H3-K9 Antibody at dilution of 1/100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) at 1/500 dilution. Blue: DAPI for nuclear staining.



Chromatin immunoprecipitation analysis of extracts of HeLa cells, using MonoMethyl-Histone H3-K9 antibody and rabbit IgG. The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.



Chromatin immunoprecipitation analysis of extracts of HeLa cells, using MonoMethyl-Histone H3-K9 antibody and rabbit IgG. The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.

Histone H3K9me1 Antibody is a Rabbit Polyclonal antibody against Histone H3K9me1. Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes.

Datasheet

Version: 6.0.0
Revision date: 02 Jun 2025



to form higher order chromatin structures. This gene is intronless and encodes a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is located separately from the other H3 genes that are in the histone gene cluster on chromosome 6p22-p21.3.

Target:	Histone H3K9me1
Clonality:	Polyclonal
Reactivity:	Human, Mouse, Rat
Tested Applications:	ELISA, WB, IHC, IF/ICC, DB, ChIP
Host:	Rabbit
Recommended dilutions:	ELISA: 1 µg/ml, WB: 1/500 - 1/1000, IHC-P: 1/50 - 1/200, IF/ICC: 1/50 - 1/200, DB: 1/500 - 1/2000, ChIP: 5 µg antibody per 5 µg - 10 µg of Chromatin, ChIP-seq: 1/20 - 1/100. Not tested in IHC-F. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	A synthetic monomethylated peptide around K9 of human histone H3.
Isotype:	IgG
Form:	Liquid
Purification:	Purified by affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	Q16695 (UniProt , ExPASy)
Gene Symbol:	HIST3H3
GeneID:	8290
NCBI Accession:	NP_003520.1
KEGG:	hsa:8290
String:	9606.ENSP00000355657
Molecular Weight:	Calculated MW: 16 kDa Observed MW: 14 kDa
Buffer:	PBS, pH 7.3, containing 0.05% Proclin-300, 50% glycerol.

Datasheet

Version: 6.0.0

Revision date: 02 Jun 2025



Concentration: > 0.2 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only