



## Tryptophan Hydroxylase 2 (TPH2) Antibody

Catalogue No.:abx005394



Western blot analysis of various lysates using TPH2 Antibody at 1/1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1/10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Exposure time: 90s.



Western blot analysis of various lysates using TPH2 Antibody at 1/1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1/10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Exposure time: 5s.

TPH2 Antibody is a Rabbit Polyclonal antibody against TPH2. This gene encodes a member of the pterin-dependent aromatic acid hydroxylase family. The encoded protein catalyzes the first and rate limiting step in the biosynthesis of serotonin, an important hormone and neurotransmitter. The human genome contains two related tryptophan hydroxylases, one on chromosome 11p15-p14 and one on chromosome 12q21. This gene is expressed predominantly in the brain stem. Mutations in this gene may be associated with psychiatric diseases such as bipolar affective disorder and major depression.

Target: Tryptophan Hydroxylase 2 (TPH2)

Clonality: Polyclonal

Reactivity: Human, Mouse, Rat

Tested Applications: ELISA, WB

Host: Rabbit

Recommended dilutions: ELISA: 1 µg/ml, WB: 1/500 - 1/1000. Optimal dilutions/concentrations should be determined by the

end user.

Conjugation: Unconjugated

**Immunogen:** Synthetic peptide corresponding to TPH2. The exact sequence is proprietary.

Isotype: IgG

## **Datasheet**

Version: 5.0.0 Revision date: 30 Jul 2025



Form: Liquid

**Purification:** Purified by affinity chromatography.

**Storage:** Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

UniProt Primary AC: Q8IWU9 (<u>UniProt</u>, <u>ExPASy</u>)

Gene Symbol: TPH2

GenelD: <u>121278</u>

NCBI Accession: NP\_775489.2

**KEGG:** hsa:121278

String: <u>9606.ENSP00000329093</u>

Molecular Weight: Calculated MW: 56 kDa

Observed MW: 56/52 kDa

**Buffer:** PBS, pH 7.3, containing 0.02% sodium azide, 50% glycerol.

**Concentration:** > 0.2 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC,

THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL

CONSUMPTION.