


Goat Anti-Rat IgG H&L Antibody (TRITC)

Catalogue No.: abx005555



Flow cytometry: 1×10^6 RK13 cells (negative control, left) and RK13-CD20 transfection cells (right) were surface-stained with rat anti-mouse CD20 Antibody (1/100, orange line) or secondary antibody only (blue line). Non-fluorescently stained RK13 and RK13 transfection cells were used as blank control (red line). Rhodamine Goat Anti-Rat IgG (H+L)(1/200) was used as a secondary antibody.

Goat Anti-Rat IgG (H+L) Antibody (TRITC) is a Goat antibody against Rat IgG (H+L). This product is for research use only, not for diagnostic or therapeutic use.

Target: Rat IgG H&L

Clonality: Polyclonal

Reactivity: Rat

Tested Applications: IF/ICC, FCM

Host: Goat

Recommended dilutions: IF/ICC: 1/50 - 1/200, FCM: 1/50 - 1/200. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: TRITC

Excitation/Emission: 552/576

Laser Line: 550

Immunogen: Rat IgG

Isotype: IgG

Form: Liquid

Purification: Purified by affinity chromatography.

Storage: Aliquot and store at -20°C . Avoid repeated freeze/thaw cycles.

Datasheet

Version: 2.0.0

Revision date: 07 Sep 2025



Buffer: PBS, pH 7.3, containing 0.025% sodium azide, 0.75% BSA, 50% glycerol.

Concentration: > 0.2 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only