



Goat Anti-Mouse IgG H&L Antibody (TRITC)

Catalogue No.: abx005559



Immunofluorescence analysis of NIH/3T3 cells using TRITC Goat Anti-Mouse IgG (H+L) at dilution of 1/200 (40x lens). Blue: DAPI for nuclear staining.



Flow cytometric analysis of Positive antibody Human Calcitonin R (2.5 µg/ml) in various cells (orange) compare to Mouse isotype control (blue) and non-staining control (Red). The secondary antibody used was TRITC Goat Anti-Mouse IgG (H+L) at 1/100.

Goat Anti-Mouse IgG (H+L) Antibody (TRITC) is a Goat antibody against Mouse IgG (H+L). This product is for research use only, not for diagnostic or therapeutic use.

Target: Mouse IgG H&L

Clonality: Polyclonal

Reactivity: Mouse

Tested Applications: IF/ICC, FCM

Host: Goat

Recommended dilutions: IF/ICC: 1/50 - 1/200, FCM: 1/50 - 1/200. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: TRITC

Excitation/Emission: 552/576

Laser Line: 550

Immunogen: Mouse IgG

Datasheet

Version: 2.0.0

Revision date: 07 Sep 2025



Isotype:	IgG
Form:	Liquid
Purification:	Purified by affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
Buffer:	PBS, pH 7.3, containing 0.025% sodium azide, 0.75% BSA, 50% glycerol.
Concentration:	> 0.2 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only