



Oxidative Stress-Induced Growth Inhibitor 2 (OSGIN2) Antibody

Catalogue No.: abx122927



Western blot analysis of recombinant protein using OSGIN2 Antibody.



Immunohistochemistry analysis of formalin-fixed paraffin-embedded foetal thyroid gland showing cytoplasmic staining with OSGIN2 Antibody (dilution 1/100).

Rabbit Polyclonal against the OSGIN2 protein.

Target:	Oxidative Stress-Induced Growth Inhibitor 2 (OSGIN2)
Clonality:	Polyclonal
Reactivity:	Human
Tested Applications:	ELISA, WB, IHC
Host:	Rabbit
Recommended dilutions:	ELISA: 1/20000 - 1/80000, WB: 1/500 - 1/2000, IHC: 1/100 - 1/200. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	Recombinant fragment corresponding to 436-632 AA of human OSGIN2.
Isotype:	IgG
Form:	Lyophilized
Purification:	Purified by antigen affinity column chromatography.

Datasheet

Version: 5.0.0

Revision date: 10 Jul 2025



Reconstitution:	Reconstitute in 100 µl of sterile distilled H ₂ O with 50% glycerol.
Storage:	Store at -20 °C. Avoid repeated freeze/thaw cycles.
NCBI Accession:	BC031054
Molecular Weight:	26 kDa (505aa, 57 kDa and 549aa, 62 kDa)
Buffer:	Prior to lyophilization: 1% BSA and 0.02% NaN ₃ .
Specificity:	Predicted to react with Mouse and Rat OSGIN2.
Concentration:	Lyophilized form: Not applicable. After reconstitution: 1 mg/ml.
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only