

Human FAF1 siRNA

Catalogue No.:abx901826

siRNA to inhibit FAF1 expression using RNA interference.

This product is provided as three 5 nmol vials (15 nmol) or 2x three 5 nmol vials (30 nmol) of lyophilized siRNA oligo duplexes. Each vial contains slightly different sequences to ensure full knockout of the gene. The duplexes can be transfected individually or pooled together to achieve knockdown of the target gene, which is most commonly assessed by qPCR or western blot.

Target:	FAF1				
Reactivity:	Human				
Tested Applications:	RNAi				
Host:	Synthetic		C	6	
Recommended	Ontimal dil	utiona/concentrati	one should be determined	t by the and upor	
	Optimal dilutions/concentrations should be determined by the end user.				
dilutions:	Plate	Final Medium	Final siRNA	20 µM siRNA	Lipofectamine 2000
	(wells)	Volume (ml)	Concentration (nM)	Volume (μl) 0.5	Volume (μl) 0.25
	96	0.1	50	0.25	0.25
	00	0.1	10	0.05	0.25
			100	2.5	1
	24	0.5	50	1.25	1
			10	0.25	1
			100	5	2
	12	1	50	2.5	2
			10	0.5	2
			100	10	5
	6	2	50	5	5
			10	1	5
Form:	Lyophilized	b			
Purity:	> 97%				
Overlite Overteele		- 4: -!			4
Quality Control:	Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure appropriate				
	coupling e	fficiency. The oligo	o is subsequently purified	by affinity-solid phas	e extraction. The
	annealed I	RNA duplex is furt	her analyzed by mass spe	ectrometry to verify th	e exact composition of
		-	ared to the previous lot by		-
		•	area to the previous lot by	mass specironieu y	
	to-lot cons	istency.			
Storage:	Shipped at	t 4 °C. Store at -20) °C for up to one year.		
UniProt Primary AC:	Q9UNN5 (<u>UniProt</u> , <u>ExPASy</u>)				
Gene Symbol:	FAF1				

Datasheet Version: 1.0.0 Revision date: 24 May 2025



GenelD:	<u>11124</u>		
NCBI Accession:	NM_007051.2		
KEGG:	hsa:11124		
Specificity:	FAF1 siRNA (Human) is a target-specific 19-23 nt siRNA oligo duplexes designed to knock down gene expression.		
Note:	This product is for research use only.		
Directions for use:	 1. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. 2. Resuspend the siRNA oligos to an appropriate concentration with DEPC water (e.g. resuspend one vial of 5 nmol siRNA oligo in 250 µl of DEPC water for a final concentration of 20 µM). 3. Transfect with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. 		