



## Activin A Receptor Type 1C (ACVR1C) Antibody


Catalogue No.: abx000813




Western blot analysis of various lysates using ACVR1C Antibody at 1/1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1/10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Exposure time: 180s.




Western blot analysis of various lysates using ACVR1C Antibody at 1/1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1/10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Exposure time: 180s.



Immunohistochemistry analysis of paraffin-embedded Rat brain using ACVR1C Antibody at dilution of 1/100 (40x lens). Microwave antigen retrieval performed in 0.01 M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human stomach using ACVR1C Antibody at dilution of 1/100 (40x lens). Microwave antigen retrieval performed in 0.01 M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse stomach using ACVR1C Antibody at dilution of 1/100 (40x lens). Microwave antigen retrieval performed in 0.01 M PBS Buffer (pH 7.2) prior to IHC staining.

# Datasheet

Version: 4.0.0  
Revision date: 10 Apr 2025



ACVR1C Antibody is a Rabbit Polyclonal antibody against ACVR1C. ACTR-IC (Activin receptor type 1C), also referred to as Activin receptor-like kinase 7 (ALK-7), is a type I serine/threonine kinase receptor. ACTA-IC contains an extracellular binding domain, an intracellular serine/threonine kinase domain preceded by a GS box and a transmembrane domain. It is expressed throughout the digestive and central nervous system and localizes to the cell surface. Four ACTR-IC transcripts are generated by alternative splicing. Transcript 1 is the functional full length receptor, transcript 2 lacks a complete receptor binding domain and transcripts 3 and 4 are soluble proteins that lack a transmembrane domain. ACTR-IC is a receptor for Activin AB, Activin B and Nodal. In pancreatic cells, ACTR-IC forms a complex with Activin receptor type IIB (ACTR-IIB). The kinase domain of ACTR-IC can induce Smad2 and Smad3 signalling pathways. In some cell lines, ACTR-IC overexpression induces apoptosis and inhibits proliferation.

<b>Target:</b>	Activin A Receptor Type 1C (ACVR1C)
<b>Clonality:</b>	Polyclonal
<b>Reactivity:</b>	Human, Mouse, Rat
<b>Tested Applications:</b>	ELISA, WB, IHC
<b>Host:</b>	Rabbit
<b>Recommended dilutions:</b>	ELISA: 1 µg/ml, WB: 1/500 - 1/1000, IHC-P: 1/100 - 1/200. Not tested in IHC-F. Optimal dilutions/concentrations should be determined by the end user.
<b>Conjugation:</b>	Unconjugated
<b>Immunogen:</b>	Recombinant fusion protein containing a sequence corresponding to amino acids 22-113 of human ACVR1C.
<b>Isotype:</b>	IgG
<b>Form:</b>	Liquid
<b>Purification:</b>	Purified by affinity chromatography.
<b>Storage:</b>	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
<b>UniProt Primary AC:</b>	Q8NER5 ( <a href="#">UniProt</a> , <a href="#">ExPASy</a> )
<b>Gene Symbol:</b>	ACVR1C
<b>GeneID:</b>	<a href="#">130399</a>
<b>NCBI Accession:</b>	NP_660302.2
<b>KEGG:</b>	hsa:130399
<b>String:</b>	<a href="#">9606.ENSP00000243349</a>

# Datasheet

Version: 4.0.0

Revision date: 10 Apr 2025



**Molecular Weight:** Calculated MW: 55 kDa  
Observed MW: 55 kDa

**Buffer:** PBS, pH 7.3, containing 0.05% Proclin-300, 50% glycerol.

**Concentration:** > 0.2 mg/ml

**Note:** THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only