



Nuclear Factor of Activated T-Cells 3 (NFATC3) Antibody

Catalogue No.: abx011684



WB analysis of Jurkat cells, using NFATC3 antibody (1/1000 dilution).



IHC-P analysis of human kidney tissue, using NFATC3 antibody (1/100 dilution).

Nuclear Factor of Activated T-Cells 3 (NFATC3) Antibody is a Rabbit Polyclonal antibody for the detection of NFATC3.

Acts as a regulator of transcriptional activation. Plays a role in the inducible expression of cytokine genes in T-cells, especially in the induction of the IL-2.

Target: Nuclear Factor of Activated T-Cells 3 (NFATC3)

Clonality: Polyclonal

Reactivity: Human

Tested Applications: ELISA, WB, IHC

Host: Rabbit

Recommended dilutions: WB: 1/200 - 1/2000, IHC: 1/50 - 1/200. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: nuclear factor of activated T-cells, cytoplasmic, calcineurin-dependent 3

Isotype: IgG

Datasheet

Version: 2.0.0
Revision date: 05 Mar 2025



Form:	Liquid
Purity:	≥ 95% (SDS-PAGE)
Purification:	Purified by immunogen affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
Validity:	12 months.
UniProt Primary AC:	Q12968 (UniProt , ExPASy)
Gene Symbol:	NFATC3
GeneID:	4775
OMIM:	602698
HGNC:	7777
KEGG:	hsa:4775
Ensembl:	ENSG00000072736
String:	9606.ENSP00000300659
Molecular Weight:	Observed MW: 130-170 kDa
Buffer:	PBS, pH 7.3, with 0.02% sodium azide and 50% glycerol.
Concentration:	2 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.