



Abbexa Ltd, Innovation Centre, Cambridge Science Park, Cambridge, CB4 0EY, UK
 Telephone: +44 (0) 1223 755950 - Fax: +44 (0) 1223 755951 - E-Mail: info@abbexa.com

ID2 Antibody


Catalogue No.: abx015888




Western blot analysis using ID2 antibody against human ID2 recombinant protein. (Expected MW is 17.3 kDa).




Flow cytometric analysis of SK-N-SH cells using ID2 antibody (green) and negative control (purple).



Immunohistochemical analysis of paraffin-embedded breast cancer tissues using ID2 antibody with DAB staining.



Immunohistochemical analysis of paraffin-embedded rectum cancer tissues using ID2 antibody with DAB staining.



Red: Control Antigen (100ng) ; Purple: Antigen (10ng) ; Green: Antigen (50ng) ; Blue: Antigen (100ng).

The protein encoded by this gene belongs to the inhibitor of DNA binding family, members of which are transcriptional regulators

Abbexa Ltd, Innovation Centre, Cambridge Science Park, Cambridge, CB4 0EY, UK
Telephone: +44 (0) 1223 755950 - Fax: +44 (0) 1223 755951 - E-Mail: info@abbexa.com

that contain a helix-loop-helix (HLH) domain but not a basic domain. Members of the inhibitor of DNA binding family inhibit the functions of basic helix-loop-helix transcription factors in a dominant-negative manner by suppressing their heterodimerization partners through the HLH domains. This protein may play a role in negatively regulating cell differentiation. A pseudogene of this gene is located on chromosome 3.

Target:	ID2
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Clonality:	Monoclonal
Tested Applications:	ELISA, WB, IHC, FCM
Recommended dilutions:	ELISA: 1/10000, WB: 1/500 - 1/2000, IHC: 1/200 - 1/1000, FCM: 1/200 - 1/400. Optimal dilutions/concentrations should be determined by the end user.
Immunogen:	Purified recombinant fragment of human ID2 (AA: 1-134) expressed in E. coli.
Isotype:	IgG ₁
Conjugation:	Unconjugated
Storage:	Aliquot and store at -20 °C. Avoid repeated freeze/thaw cycles.
Molecular Weight:	15 kDa
GeneID:	3398 15902 25587
Buffer:	Ascitic fluid containing 0.03% sodium azide.
Availability:	Shipped within 5-10 working days.
Note:	This product is for research use only.